#### Example /

Pest: Tetranychus urticae

Crop: Hops

The compound of the formula (I) (240SC) was tested at a concentration of 0.0048% a.i. in comparison with cis-cyhalothrin (050EC) at 0.005% a.i. The compound of the formula (I) was used at an early stage of the infestation of the pest, while cis-cyhalothrin was applied when the infestation had progressed to a high level.

The spray mixture (2000 I/ha) was applied using a knapsack sprayer operated with compressed air.

10 The plot size was 6 plants, the number of replications per test variant was 2.

The activity against spider mites was determined 4, 14 and 21 days (compound of the formula (I)) and 4, 11 and 18 days (cis-cyhalothrin) after the treatment by counting the live animals/leaf (10 leaves/plot) and calculating the efficacy using Abbott's formula.

#### Table

#### 15 Tetranychus urticae / Hops

Active ingredients	Concentration	Effica	cy in %	Abbott		
	% a.i./ha	4*	11*	14*	18*	21*
Compound of the formula (I) (240 SC)	0.0048 £96g/ha	90	<b></b>	94		93
Cis-cyhalothrin (050 EC)	0.005	59 * T	57 Bays afte	r treatme	o	

### Example >

Pest: Tetranyous urticae

Crop: Hops

The compound of the formula (I) (240 SC) was tested at a concentration of 0.0144% a.i. in comparison with the standard Amitraz (200 EC) at 0.05% a.i. The mixture was sprayed once.

The spray mixture (2200 l/ha) was applied using a trailed machine operated by a tractor. The plot size was 60 plants, and 2 replications were carried out per test variant.

The activity against spider mites was determined 5, 12, 19 and 34 days after the treatment by counting the live animals/leaf (60 leaves/plot) and calculating the efficacy using Abbott's formula.

#### 10 Table B

Tetranychus urticae / Hops

Active ingredients	Concentration	Eff	icacy in % A	bbott
·	% a.i./ha	5*	12*	19*
Compound of the formula (I) (240 SC)	0.0144 \$^ca:320g Qa	89.7	98.1	99.6
Amitraz (200 EC)	0.05	80.1	96.3	92.2

\* Days after treatment

### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

**Applicants** 

R. Fischer et al.

Serial No.

10/563,803

2

Filed

June 28, 2006

For

Use of 2,2-Dimethyl-3-(2,4-dichlorophenyl)-2-oxo1-

oxaspiro[4,5]dec-3-en-4-yl Butanoate for Controlling

Acarides

Group Art Unit

161

Examiner

BROOKS, K.L.

#### DECLARATION UNDER 37 C.F.R. §1.132

- I, Dr Robert Brinkmann, hereby declare:
- that I am an Agronomist having studied at the University of Hannover,
   Germany;
- that I received my doctor's degree in phytopathology at the University of Hannover, Germany in 1987;
- that I entered the employ of Bayer in 1991;
- that I am specialized in plant protection (phytopathology);
- that the following tests have been cerried out under my supervision and direction

Pest: Tarsonemus pallidus

Crop: Strawberries Variety: Selekt

The compound of the formula (I) (240SC) is tested with 210 g a i./ha in comparison with Abamectin EC (020) with 23 g a i./ha.

The spray mixture (300 l/ha) is applied with a knapsack sprayer powered with compressed air of 2 bars.

The size of the plot is 12 m<sup>2</sup> and the number of repeated applications per test variant is 3.

The activity against red spider mites is determined 5 and 12 days after treatment by counting the number of living animals per leaf and calculating the degree of activity by the Abbott method.

## Tarsonemus pallidus / Strawberries

#### **Active compounds**

The section of the se	Concentration in g a.i./ha	Degree of activit in Abbott %	
Compound of the formula (I) (240 SC)	210	5* 100	12* 100
Abamectin EC (020)	23	100	100

\*days after treatment

The undersigned declarant hereby declares that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

15.7.204

Date

XX (Name)

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants

R. Fischer et al.

Serial No.

10/563,803

Filed

June 28, 2006

For

Use of 2,2-Dimethyl-3-(2,4-dichlorophenyl)-2-oxo1-

oxaspiro[4,6]dec-3-en-4-yl Butanoate for Controlling Acarides

Group Art Unit

161

Examiner

BROOKS, K. L.

## DECLARATION UNDER 37 C.F.R. § 1.132

- I, John Bell, hereby declare:
- that I am a Pest Management Specialist having studied at Mississippi State
  University, USA;
- that I received my Master of Agriculture degree at Mississippi State University,
   USA in 1981;
- that I entered the employ of Bayer in 1982;
- that I am specialized in plant protection (phytopharmacology);
- that the following tests have been carried out under my supervision and direction

5

Plots of around 32 m<sup>2</sup> with cotton of the cultivar "Maxxa" are treated in four replications against Tetranychus urticae. Application takes place using a tractor-operated sprayer bar, with a pressure of 50 PSI. The active ingredient of Example (I) (240 SC) is tested against the commercial standard Zephyr (BC 0.15) at the stated application rates. The water application rate is 350 l/ha.

Evaluation is made 7, 14 and 21 days after treatment, by scoring the destruction of the animals on the plants.

Active ingredient	Application rate (g) a.i./ha	Activity (% Abbott)		
ingroutone	annual annual state of the Annual	7 d	14 d	21 d
Zephyr	TI	31	73	89
Example (I)	150	0	15	58

5

Plots of around 12 m<sup>2</sup> with cotton of the cultivar "Acala GC510" are treated in three replications against Tetranychus urticae. Application takes place using a tractor-operated sprayer bar, with a pressure of 60 PSI. The active ingredient of Example (I) (200 SC) is tested against the commercial standard Zephyr (BC 0.15) at the stated application rates. The water application rate is 500 l/ha.

Evaluation is made 9 days after treatment, by scoring the destruction of the animals on the plants.

Active	Application rate (g) a.i./ha	Activity (% Abbott)
ingredient		
	CONTRACTOR AND A SECOND OF THE SECOND	b 6
Example (I)	125	60
Zephyr	11.	93

5

Plots of around 32 m<sup>2</sup> with cotton of the cultivar "Maxxa" are treated in six replications against Tetranychus urticae. Application takes place using a tractor-operated sprayer bar, with a pressure of 50 PSI. The active ingredient of Example (I) (240 SC) is tested against the commercial standard Zephyr (EC 0.15) at the stated application rates. There are two applications, made 10 days apart. The water application rate is 30 US gallons ≈ about 132 1/ha.

Bualuation is made 7 days after the first treatment (1.B) and 7 and 14 days after the second treatment (2.B), by scoring the destruction of the animals on the plants.

Active	Application rate (g) a.i./ha	Activity (% Abbott)		
ingredient				
	майшай көлөмөлөөнөө үнүүнүү дүйнө сүчт чэгээ үзэх эрэхэ (А) уйй А А уу хэрэх дүйдөө байлай байлай байлай байла	7 d-1.B	7 d-2.B	14 d-2.B
Zephyr	okaning processor and a single-part of the contract of the con	69	96	98
Example (I)	125	74	79	84

Plots of around 10.5 m<sup>2</sup> with cotton of the cultivar "Maxxa" are treated in four replications against Tetranychus urticae. Application takes place using a compressed-air operated sprayer bar, with a pressure of 2 atm. The active ingredient of Example (I) (240 SC) is tested against the commercial standard Zephyr (EC 0.15) at the stated application rates. The water application rate is 350 l/ha.

5 Evaluation is made 14 and 21 days after treatment, by scoring the destruction of the animals on the plants.

Active	Application rate (g) a.i./ha	Activity (9	6 Abbott)
ingredient			
	हुं होता कार्यक्रमाता विकास कर कर कार्यक्रमाता विकास प्रकार के प्रकार कार्यक्रमा है कि विकास कार्यक्रमा के प्र	14 d	21 d
Zepliyr	× • • • • • • • • • • • • • • • • • • •	80	89
Example (I)	330	20	44

## Example S

5.

Plots of around 28 m<sup>2</sup> with cotton of the cultivar "Maxxa" are treated in four replications against Tetranychus urticae. Application takes place using a tractor-operated sprayer bar, with a pressure of 2.72 atm. The active ingredient of Example (I) (240 SC) is tested against the commercial standard Zephyr (EC 0.15) at the stated application rates. The water application rate is 350 l/ha.

Evaluation is made 14 days after treatment, by scoring the destruction of the animals on the plants.

Active	Application rate (g) a.i./ha	Activity (% Abbott)
ingredient		
±- ••• • • • • • • • • • • • • • • • • •	о том от	14 d
Zephyr	11	83
Example (I)	330	30

The undersigned declarant hereby declares that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

July 5, 4011

John Rail